

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets

(11) Publication number:

0 318 029  
A2

(12)

## EUROPEAN PATENT APPLICATION

(21) Application number: 88119675.2

(22) Date of filing: 25.11.88

(51) Int. Cl. 4: C07D 211/14 , C07D 211/32 ,  
C07D 211/40 , C07D 401/12 ,  
C07D 211/18 , C07D 211/58 ,  
C07D 223/04 , A61K 31/445 ,  
A61K 31/55

(30) Priority: 27.11.87 JP 299438/87  
28.03.88 JP 74127/88

(43) Date of publication of application:  
31.05.89 Bulletin 89/22

(54) Designated Contracting States:  
AT BE CH DE FR GB IT LI LU NL SE

(71) Applicant: Eisai Co., Ltd.  
6-10, Kolshikawa 4-chome Bunkyo-ku  
Tokyo 112(JP)

(72) Inventor: Sugimoto, Hachiro  
3073-13, Kashiwadamachi  
Ushiku-shi Ibaraki(JP)  
Inventor: Nakamura, Takaharu  
4-9, Tsukushino 2-chome  
Abiko-shi Chiba(JP)  
Inventor: Karibe, Norio  
Shinzanryo 19-13, Kasuga 4-chome  
Tsukuba-shi Ibaraki(JP)  
Inventor: Saito, Isao  
670-63, Shimohirooka  
Tsukuba-shi Ibaraki(JP)  
Inventor: Higurashi, Kunizo  
4-3, Midori 2-chome  
Sumida-ku Tokyo(JP)


Inventor: Yonaga, Masahiro  
901-405, 6-2, Takezono 1-chome  
Tsukuba-shi Ibaraki(JP)  
Inventor: Kaneko, Takeru  
6-1, 17, Matsushiro 2-chome  
Tsukuba-shi Ibaraki(JP)  
Inventor: Nakazawa, Takahiro  
1-810, 531, Ohaza Miyawada Fujishiromachi  
Kitasouma-gun Ibaraki(JP)  
Inventor: Ueno, Masataka  
Parkside Doho 205 17-10, Ninomiya 2-chome  
Tsukuba-shi Ibaraki(JP)  
Inventor: Yamatsu, Kiyomi  
23-7, Imazumidai 7-chome  
Kamakura-shi Kanagawa(JP)  
Inventor: Ueno, Kohshi  
Corpo Namikicho 102 22-7, Namiki 3-chome  
Tsukuba-shi Ibaraki(JP)  
Inventor: Ikeda, Masuhiro  
6-6, Toukoudai 3-chome  
Tsukuba-shi Ibaraki(JP)

(74) Representative: Hansen, Bernd, Dr.rer.nat. et  
al  
Hoffmann, Eitle & Partner Patentanwälte  
Arabellastrasse 4 Postfach 81 04 20  
D-8000 München 81(DE)

EP 0 318 029 A2

(54) Cyclic amine and pharmacological composition.

(57) A cyclic amine is defined by the formula:



( I )

**EP 0 318 029 A2**

in which A is naphthyl, phenyl, quinolyl or alkyl, X is a group containing carbonyl, n is 2 or 3 and B is alkyl, halogen, phenyl or -YZ, Y being -O-, -CO- or -NH-, Z being phenyl, naphthyl, pyridyl, alkyl or cycloalkyl. It is useful for the making of a medicament to treat a mental symptom associated with a cerebrovascular disease and respective pharmaceutical compositions are disclosed.

## Cyclic Amine and Pharmacological Composition

The present invention relates to a novel cyclic amine derivative which exhibits an excellent activity as a pharmaceutical.

## 5 [Background of the Invention and Prior Art]

Various attempts have been made to treat cerebrovascular diseases with a drug. For example, cerebrovasodilators and cerebral metabolism activators are used. So far, however, there has been no drug which is very useful for the treatment of the cerebrovascular diseases. In particular, with respect to  
10 cerebrovascular dementia, mental dysfunction, etc. among various symptoms accompanying cerebrovascular diseases, no effective drug has yet been developed.


However, in recent years, a theory that an excitatory amino acid liberated after ischemia seriously participates in the death of nerve cells has attracted worldwide attention. This theory was substantiated by the facts that ischemia brings about liberation of a glutamate and the cell injury derived from ischemia is  
15 antagonized by a glutamate antagonist.

The present inventors have paid their attention to the neurotoxin activity of the above-described glutamate in starting the search for a new agent for treating cerebrovascular diseases, particularly an agent for improving a mental symptom accompanying the cerebrovascular diseases and have repeated studies on various compounds for a long period of time. As a result, we have found that a cyclic amine derivative  
20 which will be described hereinbelow exhibits an excellent activity on the suppression of glutamate liberation, which have led to the completion of the present invention.

Accordingly, an object of the present invention is to provide a cyclic amine derivative or pharmacologically acceptable salts thereof effective as an agent for improving mental symptoms accompanying cerebrovascular diseases, such as cerebral apoplexy, cerebral hemorrhage, cerebral infarction and cerebral  
25 arteriosclerosis, and polyinfarctive dementia, and a process for preparing said compound and pharmacologically acceptable salts thereof. Another object of the present invention is to provide a pharmaceutical comprising as an effective ingredient said compound or pharmacologically acceptable salts thereof.

## 30 [ Summary of the Invention ]


The object compound of the present invention is a cyclic amine and pharmacologically acceptable salts thereof represented by the following general formula (1):




wherein A is a naphthyl group or a naphthyl group substituted with a lower alkoxy or hydroxy; phenyl group or a phenyl group substituted with a lower alkyl group or a halogen; a quinolyl group; or a lower alkyl group, n is an integer of 2 or 3, X is a group represented by the formula

45  $\text{O} \parallel$   
-C-CH<sub>2</sub>-, a group of the formula

$\text{O} \parallel$   
-C-, a group represented by the formula



or a group represented by the formula



- 5 B is a lower alkyl group; a phenyl group which may be substituted with 1 to 3 same or different substituents selected from the group consisting of a halogen, a lower alkyl group, and a lower alkoxy group; or a group represented by the formula -Y-Z wherein Y is a group represented by the formula -O-, a group represented by the formula



- 10 Z is a phenyl group which may be substituted with one or two same or different substituents selected from the group consisting of a halogen a lower alkyl group, and a lower alkoxy group; a naphthyl group; a pyridyl group; a lower alkyl group; or a cycloalkyl group.

The term "lower alkyl group" used in the above definition of A, B, and Z with respect to the compound (I) of the present invention is intended to mean a straight-chain or branched alkyl group having 1 to 6 carbon atoms, and examples thereof include methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl (amyl), isopentyl, neopentyl, tert-pentyl, 1-methylbutyl, 2-methylbutyl, 1,2-dimethylpropyl, hexyl, isohexyl, 1-methylpentyl, 2-methylpentyl, 3-methylpentyl, 1,1-dimethylbutyl, 1,2-dimethylbutyl, 2,2-dimethylbutyl, 1,3-dimethylbutyl, 2,3-dimethylbutyl, 3,3-dimethylbutyl, 1-ethylbutyl, 2-ethyl butyl, 1,1,2-trimethylpropyl, 1,2,2-trimethylpropyl, 1-ethyl-1-methylpropyl, and 1-ethyl-2-methylpropyl groups. Among them, methyl, ethyl, propyl, isopropyl groups etc. are preferable. A methyl group is most preferable.

n is intended to mean an integer of 2 or 3. When n is 2, the compound is a piperidine derivative. When n is 3, the compound is a homopiperidine derivative.


The term "lower alkoxy group" used in the definition of A, B, and Z is intended to mean an alkoxy group derived from the above-described lower alkyl group, and preferable examples thereof include methoxy, ethoxy, propoxy, isopropoxy, butoxy, and isobutoxy groups. Among them, a methoxy group is most preferable.

Examples of the halogen atom involved in the definition of A, B, and Z include chlorine, bromine, fluorine, and iodine.

The term "pharmacologically acceptable salt" is intended to mean a commonly used nontoxic salt, and examples thereof include those of inorganic acids, such as hydrochloride, hydrobromide, sulfate and phosphate, those of organic acids, such as acetate, maleate, tartrate, methanesulfonate, benzenesulfonate, and toluenesulfonate, and salts with amino acids such as arginine, and aspartic acid.

In preferable embodiments of the cyclic amine and pharmacologically acceptable salts thereof as defined above, (1) B is a lower alkyl; (2) B is -Y-Z in which Y is -O-; (3) A is a naphthyl group or a naphthyl group having a substituent of a lower alkyl or hydroxy, X is -CO-CH<sub>2</sub>-, n is 2 and B is -Y-Z in which Y is -O-, -CO- or -NH- and Z is phenyl or phenyl having one to three substituents, either the same as or different from each other, selected from the group consisting of a halogen, a lower alkyl, a lower alkoxy and naphthyl; or (4) n is 2, B is a lower alkyl or -YZ and Z is defined as above however to exclude cycloalkyl.


The invention provides a process for preparing a cyclic amine or a pharmacologically acceptable salt thereof as defined above, which comprises the step of reacting a halide compound having the formula. A-X-Hal in which Hal is a halogen and A and X each are defined above, with a compound having the formula:



50 in which n and B each are defined above.

The invention also provides a pharmacological composition which comprises a pharmacologically effective amount of a cyclic amine or a pharmacologically acceptable salt thereof as defined above and a pharmacologically acceptable carrier and then a method for improving, treating or preventing a mental symptom associated with a cerebrovascular disease, which comprises administering a pharmacologically effective amount of a cyclic amine or a pharmacologically acceptable salt thereof as defined above to a subject suffering from the mental symptom associated with a cerebrovascular disease.

The compounds of the present invention may be prepared by various processes. A representative process for preparing the compounds of the present invention will now be described.

Process of preparation

25 wherein A, X, B, and n are as defined above and Hal is a halogen atom.

Specifically, a halide represented by the general formula (II) is reacted with a compound represented by the general formula (III) to prepare a compound (I) which is an object compound of the present invention.

This reaction is dehydrohalogenation conducted by any ordinary method while heating in the absence of any solvent or in an organic solvent which does not participate in the reaction, i.e., one selected from among alcoholic solvents, such as methanol, ethanol, and butanol, benzene, toluene, xylene, tetrahydrofuran, chloroform, carbon tetrachloride, and dimethylformamide. In this case, favorable results can be attained by conducting the reaction in the presence of an inorganic salt, such as sodium hydrogencarbonate, potassium carbonate, sodium carbonate or caustic soda, or an organic base such as triethylamine, pyridine, pyrimidine or diethylaniline.

35 The effect of the present invention will now be described in more detail with reference to the following pharmacological experiment.

Experimental ExampleGlutamate liberation inhibitory activity in cerebral cortex slice

45 A male SD rat was decapitated, and cerebral cortices on both sides were then extirpated to prepare a slice having a thickness of 750  $\mu$ m (weight: 10 to 20 mg) with a McIlwain tissue chopper. The slice was incubated in a Krebs-Henseleit liquid containing a test compound in the form of a solution thereof for 30 minutes. Then the slice was incubated in one ml of a Krebs-Henseleit liquid containing 50 mM of KCl and the test compound solution for 60 mins. For control, a solvent was added to a Krebs-Henseleit liquid containing only 50 mM of KCl.

50 The glutamate liberated in the solution was determined by HPLC, and the glutamate liberation inhibitory activity (%) in  $10^{-4}$  M of each test compound solution was calculated by the following equation:

$$(1 - \frac{\text{glutamate in each test compound solution}}{\text{amount of glutamate in control group}}) \times 100$$

Table 1:

glutamate liberation inhibitory activity	
test compound	glutamate liberation inhibitory activity (%) $1 \times 10^{-4}$ M
compd. of Ex. 4	46
compd. of Ex. 6	42
compd. of Ex. 10	49
compd. of Ex. 12	40
compd. of Ex. 16	44
compd. of Ex. 17	39
compd. of Ex. 18	10
compd. of Ex. 21	16

It is apparent from the above-described pharmacological experiment that the compounds of the present invention have a remarkable glutamate liberation inhibitory activity and exhibit a cell injury protective action and a learning disturbance improving action after ischemia based on this activity. Therefore, the compounds of the present invention exhibit a useful pharmacological action, particularly a remarkable ischemic cerebrovascular disease improving action for the central nervous system and are useful as an agent for improving, treating or preventing mental symptoms associated with cerebrovascular diseases such as cerebral apoplexy, cerebral hemorrhage, cerebral infarction, cerebral arteriosclerosis, and various types of dementia such as polyinfarcted dementia.

Further, the compounds of the present invention exhibited high safety in toxicity tests on rats. Therefore, the present invention is valuable from this viewpoint as well.

When the compounds of the present invention are used as a pharmaceutical for the above-described diseases, they may be orally or parenterally administered. The dose will vary depending on the severity of symptom, age, sex, weight and sensitivity of patients; method of administration; time and intervals of administration and properties, formulation, and kind of pharmaceutical preparations; kind of effective ingredients, etc., so that there is no particular limitation with respect to the dose. Normally, the compounds may be administered in a dose of about 0.1 to 300 mg, preferably about 1 to 100 mg, per day per adult, ordinarily in one to four portions.

The compounds of the present invention are converted into pharmaceutical preparations in the dosage form of, e.g., injections, suppositories, sublingual tablets, and capsules according to any method which is commonly accepted in the art.

In preparing injections, the effective ingredient is blended, if necessary, with a pH modifier, a buffer, a suspending agent, a solubilizing agent, a stabilizer, a tonicity agent, a preservative, etc., followed by preparation of an intravenous, subcutaneous, or intramuscular injection according to an ordinary method. In this case, if necessary, it is possible to lyophilize these preparations according to any ordinary method.

Examples of the suspending agent include methylcellulose, Polysorbate 80, hydroxyethylcellulose, gum arabic, powdered tragacanth, sodium carboxymethylcellulose, and polyoxyethylene sorbitan monolaurate.


Examples of the solubilizing agent include polyoxyethylene hydrogenated castor oil, Polysorbate 80, nicotinamide, polyoxyethylene sorbitan monolaurate, Macrogol, and ethyl esters of castor oil fatty acids.

Examples of the stabilizer include sodium sulfite, sodium metabisulfite, and ether, and examples of the preservative include methyl p-hydroxybenzoate, ethyl p-hydroxybenzoate, sorbic acid, phenol, cresol, and chlorocresol.

#### [Examples]

Representative examples of the present invention will now be described for the purpose of aiding the understanding of the present invention. It is needless to say that the present invention will not be limited to these examples only.

## Example 1


2-(4-Methylpiperidinyl)-2'-acetonaphthone hydrochloride

2.5 g of 2-bromo-2'-acetonaphthone, 1 g of methylpiperidine, 0.1 g of potassium iodide, and 3.1 g of triethylamine were added to 100 ml of ethanol. The mixture was refluxed for 20 min. The solvent was distilled off, and dichloromethane was added to the residue. The mixture was washed with water and dried. The dichloromethane was distilled off, and the residue was purified by column chromatography (silica gel) to prepare an oleaginous and oily intended compound. The compound was converted into a hydrochloride by an ordinary method and then recrystallized to prepare 1.87 g of an intended hydrochloride.

• melting point: 202 - 203.5 °C

. elementary analysis: C <sub>18</sub> H <sub>21</sub> NO·HCl				
		C	H	N
calculated (%)	:	71.16	7.30	4.61
found (%)	:	71.21	7.59	4.50

## Example 2


2-[4-(p-Fluorobenzoyl)piperidinyl]-1'-methoxy-2'-acetonaphthone hydrochloride

1.5 g of 2-bromo-1'-methoxy-2'-acetonaphthone, 1.3 g of 4-(p-fluorobenzoyl)piperidine hydrochloride, and 1.6 g of triethylamine were added to 60 ml of ethanol. The mixture was refluxed for 15 min. The solvent was distilled off, and dichloromethane was added to the residue. The mixture was washed with water and dried. The dichloromethane was distilled off, and the residue was purified by column chromatography (silica gel) to prepare an oleaginous intended compound. The compound was converted into a hydrochloride by an ordinary method and then recrystallized to prepare 1.38 g of an intended hydrochloride.

• melting point: 195.5 - 196.2 °C

. elementary analysis: C <sub>25</sub> H <sub>24</sub> NO <sub>3</sub> F•HCl				
		C	H	N
calculated (%)	:	67.95	5.70	3.17
found (%)	:	67.88	5.64	3.02

## 10 Example 3

15 4-(p-Fluorobenzoyl)-1-(2'-naphthoyl)piperidine


25 3.8 g of 2-naphthoyl chloride, 5.5 g of 4-(p-fluorobenzoyl)piperidine hydrochloride, and 5 g of triethylamine were added to 100 ml of tetrahydrofuran. The mixture was stirred at room temperature for 1 hr. The solvent was filtered off and then distilled off. Dichloromethane was added to the residue. The mixture was washed with water and dried. The dichloromethane was distilled off, and the residue was purified by column chromatography (silica gel) to prepare a crystal. The crystal was recrystallized to prepare 5 of an intended crystal.

30 • melting point: 130.2 - 130.7 °C

. elementary analysis: C <sub>23</sub> H <sub>20</sub> NO <sub>2</sub> F				
		C	H	N
calculated (%)	:	76.44	5.78	3.88
found (%)	:	76.52	5.95	3.71

40

## Example 4

45 2-[4-(m-Nitrophenoxy)piperidyl]-2'-acetonaphthone hydrochloride

50 0.63 g of 2-bromo-2'-acetonaphthone, 0.65 g of 4-(m-nitrophenoxy)piperidine hydrochloride, and 1 g of triethylamine were added to 100 ml of ethanol. The mixture was refluxed for 30 min. The solvent was distilled off, and dichloromethane was added to the residue. The mixture was washed with water and dried. The dichloromethane was distilled off, and the residue was purified by column chromatography (silica gel) to prepare an oleaginous intended compound. The compound was converted into a hydrochloride by an



ordinary method and then recrystallized to prepare 0.15 g of an intended hydrochloride.

- melting point: 214 - 215 °C

5

. elementary analysis: $C_{23}H_{22}N_2O_4 \cdot HCl$				
		C	H	N
calculated (%)	:	64.71	5.43	6.56
found (%)	:	64.57	5.34	6.52

10

#### Examples 5 to 25

15 Various compounds shown in Table 2 were synthesized in the same manner as that of Examples 1 to 4.

20

25

30

35

40

45

50

55

Table 2

Ex. No.	structural formula	m.p. (°C)	chemical formula	elem. anal. calcd. (%) found (%)		
				C	H	N
5		125~126	$C_{15}H_{12}N_2O \cdot 2HCl$	63.01 62.90	5.77 5.52	6.68 6.83
6		201~201.5	$C_{15}H_{12}N_2O \cdot HCl$	60.77 60.01	6.60 6.79	4.22 4.19
7		206.5~207	$C_{15}H_{12}N_2O \cdot P \cdot HCl$	67.11 67.23	6.17 6.46	3.73 3.62
8		205~205.5	$C_{15}H_{12}N_2O \cdot P \cdot HCl$	60.10 60.35	6.39 6.27	4.67 4.67
9		237~238	$C_{15}H_{12}N_2O \cdot Cl \cdot P \cdot HCl$	60.46 60.60	5.33 5.32	3.53 3.48

Table 2 (continued)

Ex. No.	structural formula	m.p. (°C)	chemical formula	elem. anal. calcd. (%) found (%)		
				C	H	N
10		224.5~225	$C_{17}H_{15}ClF_2N_2O$	69.08 68.95	5.80 5.76	3.50 3.26
11		240.8~241.2	$C_{17}H_{15}ClF_2N_2O$	67.95 68.05	5.70 5.71	3.17 3.06
12		244.5~246	$C_{17}H_{15}ClF_2N_2O$	70.50 70.47	5.92 5.80	3.29 3.35
13		247.5~248	$C_{17}H_{15}ClF_2N_2O \cdot 2HCl \cdot H_2O$	59.11 59.16	5.39 5.35	5.99 5.78
14		225.5~227	$C_{17}H_{15}ClF_2N_2O$	70.50 70.43	5.92 6.05	3.29 3.12

Table 2 (continued)

Ex. No.	structural formula	m.p. (°C)	chemical formula	elem. anal. calcd. (%) found (%)		
				C	H	N
15		245~246	$C_{21}H_{19}NO_2F \cdot HCl$	67.05 67.31	5.16 5.36	3.26 3.16
16		235.5~236	$C_{21}H_{19}NO_2F \cdot HCl$	69.98 69.44	5.63 5.65	3.40 3.27
17		176.5~178	$C_{21}H_{17}Cl_2NO_2 \cdot HCl$	67.00 67.21	6.25 6.18	2.89 2.78
18		240~240.5	$C_{23}H_{19}NO_2 \cdot HCl$	75.75 75.63	5.90 6.05	3.15 3.11
19		117~118	$C_{21}H_{19}NO_2F$	76.78 76.82	5.91 6.17	3.73 3.53


Table 2 (continued)

Ex. No.	structural formula	m.p. (°C)	chemical formula	elem. anal. calcd. (%) found (%)		
				C	H	N
20		246.5~248	$C_{21}H_{19}NO_2 \cdot HCl$	71.96 72.31	6.04 6.13	3.65 3.58
21		224.5~226	$C_{21}H_{18}ClNO_2 \cdot HCl$	70.83 71.02	6.18 6.21	3.30 3.34
22		224.5~225.5	$C_{21}H_{18}ClNO_2 \cdot HCl$	71.63 71.71	6.01 6.02	3.21 3.33
23		223~244	$C_{24}H_{25}NO_2 \cdot HCl$	72.07 71.99	7.56 7.52	3.50 3.44
24		253.5~254	$C_{21}H_{19}NO_2 \cdot 2HCl$	63.45 63.17	5.79 5.65	6.43 6.20



## Claims



1. A cyclic amine and pharmacologically acceptable salts thereof represented by the following general formula:




wherein A is a naphthyl group or a naphthyl group substituted with a lower alkoxy or hydroxy; phenyl or a phenyl group substituted with a lower alkyl group or a halogen; a quinolyl group; or a lower alkyl group.

n is an integer of 2 or 3,

X is a group represented by the formula



or a group represented by the formula



B is a lower alkyl group; a phenyl group which may be substituted with 1 to 3 same or different substituents selected from the group consisting of a halogen, a lower alkyl group, and a lower alkoxy group; or a group represented by the formula -Y-Z wherein Y is a group represented by the formula -O-, a group represented by the formula



Z is a phenyl group which may be substituted with one or two same or different substituents selected from the group consisting of a halogen, a lower alkyl group, and a lower alkoxy group; a naphthyl group; a pyridyl group; a lower alkyl group; or a cycloalkyl group.


2. A cyclic amine and pharmacologically acceptable salts thereof as claimed in Claim 1, in which B is a lower alkyl.

3. A cyclic amine and pharmacologically acceptable salts thereof as claimed in Claim 1, in which B is -Y-Z in which Y is -O-.

4. A cyclic amine and pharmacologically acceptable salts thereof as claimed in Claim 1, in which A is a naphthyl group or a naphthyl group having a substituent of a lower alkyl or hydroxy, X is -CO-CH<sub>2</sub>-, n is 2 and B is -Y-Z in which Y is -O-, -CO- or -NH- and Z is phenyl or phenyl having one to three substituents, either the same as or different from each other, selected from the group consisting of a halogen, a lower alkyl, a lower alkoxy and naphthyl.

5. A cyclic amine and pharmacologically acceptable salts thereof as claimed in Claim 1, in which n is 2, B is a lower alkyl or -YZ and Z is defined in Claim 1 except for the cycloalkyl.

6. A process for preparing a cyclic amine or a pharmacologically acceptable salt thereof as defined in Claim 1, which comprises the step of reacting a halide compound having the formula: A-X-Hal in which Hal is a halogen and A and X each are defined in Claim 1, with a compound having the formula:



in which n and B each are defined in Claim 1.

7. A pharmacological composition which comprises a pharmacologically effective amount of a cyclic amine or a pharmacologically acceptable salt thereof as defined in Claim 1 and a pharmacologically acceptable carrier.

8. The use of a pharmacologically effective amount of a cyclic amine or a pharmacologically acceptable salt thereof as defined in Claim 1. for the making of a medicament for improving, treating or preventing a mental symptom associated with a cerebrovascular disease.



12

# EUROPEAN PATENT APPLICATION

21 Application number: 88119675.2

22 Date of filing: 25.11.88

51 Int. Cl.<sup>5</sup>: C07D 211/14, C07D 211/32,  
 C07D 211/40, C07D 401/12,  
 C07D 211/18, C07D 211/58,  
 C07D 223/04, A61K 31/445,  
 A61K 31/55

30 Priority: 27.11.87 JP 299438/87  
 28.03.88 JP 74127/88

43 Date of publication of application:  
 31.05.89 Bulletin 89/22.

64 Designated Contracting States:  
 AT BE CH DE FR GB IT LI LU NL SE

68 Date of deferred publication of the search report:  
 26.09.90 Bulletin 90/39

71 Applicant: Eisai Co., Ltd.  
 6-10, Koishikawa 4-chome Bunkyo-ku  
 Tokyo(JP)

72 Inventor: Sugimoto, Hachiro  
 3073-13, Kashiwadamachi  
 Ushiku-shi Ibaraki(JP)  
 Inventor: Nakamura, Takaharu  
 4-9, Tsukushino 2-chome  
 Abiko-shi Chiba(JP)  
 Inventor: Karibe, Norio  
 Shinzanryo 19-13, Kasuga 4-chome  
 Tsukuba-shi Ibaraki(JP)  
 Inventor: Saito, Isao  
 670-63, Shimohirooka  
 Tsukuba-shi Ibaraki(JP)  
 Inventor: Higurashi, Kunizo


4-3, Midori 2-chome  
 Sumida-ku Tokyo(JP)  
 Inventor: Yonaga, Masahiro  
 901-405, 6-2, Takezono 1-chome  
 Tsukuba-shi Ibaraki(JP)  
 Inventor: Kaneko, Takeru  
 6-1, 17, Matsushiro 2-chome  
 Tsukuba-shi Ibaraki(JP)  
 Inventor: Nakazawa, Takahiro  
 1-810, 531, Ohaza Miyawada Fujishinomachi  
 Kitasouma-gun Ibaraki(JP)  
 Inventor: Ueno, Masataka  
 Parkside Doho 205 17-10, Ninomiya 2-chome  
 Tsukuba-shi Ibaraki(JP)  
 Inventor: Yamatsu, Kiyomi  
 23-7, Imazumidai 7-chome  
 Kamakura-shi Kanagawa(JP)  
 Inventor: Ueno, Kohshi  
 Corpo Namikicho 102 22-7, Namiki 3-chome  
 Tsukuba-shi Ibaraki(JP)  
 Inventor: Ikeda, Masuhiro  
 6-6, Toukoudai 3-chome  
 Tsukuba-shi Ibaraki(JP)

74 Representative: Hansen, Bernd, Dr.rer.nat. et  
 al  
 Hoffmann, Eitle & Partner Patentanwälte  
 Arabellastrasse 4 Postfach 81 04 20  
 D-8000 München 81(DE)

EP 0 318 029 A3

54 Cyclic amine and pharmacological composition.

57 A cyclic amine is defined by the formula:



in which A is naphthyl, phenyl, quinolyl or alkyl, X is a group containing carbonyl, n is 2 or 3 and B is alkyl, halogen, phenyl or -YZ, Y being -O-, -CO- or -NH-, Z being phenyl, naphthyl, pyridyl, alkyl or cycloalkyl. It is useful for the making of a medicament to treat a mental symptom associated with a cerebrovascular disease and respective pharmaceutical compositions are disclosed.



European Patent  
Office

# EUROPEAN SEARCH REPORT

Application Number

EP 88 11 9675

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
X	CHEMICAL ABSTRACTS, vol. 106, no. 25, 22nd June 1987, page 645, abstract no. 213767z, Columbus, Ohio, US; & JP-A-61 227 565 (EISAI CO., LTD) 09-10-1986 * Entire abstract *	1	C 07 D 211/14 C 07 D 211/32 C 07 D 211/40 C 07 D 401/12 C 07 D 211/18 C 07 D 211/58 C 07 D 223/04 A 61 K 31/445 A 61 K 31/55
X,Y	US-A-3 576 810 (R.L. DUNCAN, G.C. HELSLEY) * Entire document; especially example 8; claim 1 *	1	
X	FR-A-2 227 868 (ROBINS) * Entire document; especially examples 1-6,9-11; claims *	1	
Y	US-A-4 075 346 (SASAJIMA et al.) * Abstract; examples; claim 1 *	1	
X	ARZNEIMITTELFORSCHUNG, vol. 24, no. 4a, 1974, pages 584-600; R. HILTMANN et al.: "2-Acylaminopyridin-Derivate mit morphinagonistischer und -antagonistischer Wirksamkeit" * Examples of table 1, page 586 *	1,2,5	TECHNICAL FIELDS SEARCHED (Int. Cl.4)  C 07 D 211/00 C 07 D 401/00 C 07 D 223/00
X	JOURNAL ORGANIC CHEMISTRY, vol. 33, no. 9, September 1968, pages 3627-3632; R.A. JOHNSON: "Conformations of alkylpiperidine amides" * Examples table 1, page 3628 *	1,2,5	
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 03-07-1990	Examiner KISSLER B.E.
<p><b>CATEGORY OF CITED DOCUMENTS</b></p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons &amp; : member of the same patent family, corresponding document</p>			



DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
X	JOURNAL OF ORGANIC CHEMISTRY, vol. 31, no. 10, October 1966, pages 3337-3342; D.M. LYNCH et al.: "Pyridylmethylnaphthalene and pyridylmethylindan derivatives" * 1-acetyl-4-methylpiperidine on page 3342 *	1,2,5	
X	JOURNAL OF MEDICINAL CHEMISTRY, vol. 13, no. 1, January 1970, pages 1-6, American Chemical Society; R.L. DUNCAN Jr. et al.: "Aroylpiperidines and pyrrolidines. A new class of potent central nervous system depressants" * Entire document; especially table 1; page 2, compounds 1,3,5 *	1-8	
L	ARZNEIMITTELFORSCHUNG, vol. 39, no. 4, 1989, pages 445-450; T. KANEKO et al.: "Anti-ischemic effect of the novel anti-ischemic agent 2-(4-(p-fluorobenzoyl)-piperidin-1-yl)-2'-acetonaphthone hydrochloride and possible mechanism of its action" * Entire document; especially structure of E-2001 in introduction; see ref. given in footnote 1 *	1-8	TECHNICAL FIELDS SEARCHED (Int. Cl.4)
X,O P	THE JAPANESE-UNITED STATES CONGRESS OF PHARMACEUTICAL SCIENCES, POSTER, abstracts, Honolulu, 2nd-7th December 1987, page 159; N. KARIBE et al.: "Anti-ischemic effect of the novel anti-ischemic agent 2-(4(p-fluorobenzoyl)-piperidin-1-yl)-2'-acetonaphthone hydrochloride and possible mechanism of its action"	1-8	
The present search report has been drawn up for all claims			
Place of search THE HAGUE		Date of completion of the search 03-07-1990	Examiner KISSLER B.E.
<b>CATEGORY OF CITED DOCUMENTS</b> X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document			